



State of New Jersey
Department of Human Services
Division of Medical Assistance & Health Services

NEWSLETTER

Volume 28 No. 05

May 2018

TO: Physicians, Advanced Practice Nurses, FQHCs, Independent Clinical Laboratories, Independent Clinics, Nurse Midwives, Medical Suppliers, and Health Maintenance Organizations – **FOR ACTION**

SUBJECT: **Revised - Preventive Care Services Guidelines and Modifier “33” (Affordable Care Act Section 4106)**

EFFECTIVE: Certain healthcare claims with service dates on or after April 1, 2018.

PURPOSE: To notify Providers about the specified preventive care services assigned a grade of A and B by the United States Preventive Services Task Force (USPSTF), and approved vaccines and their administration, recommended by the Advisory Committee on Immunization Practices (ACIP). The USPSTF A and B Recommendations may be found at: <https://www.uspreventiveservicestaskforce.org/Page/Name/uspstf-a-and-b-recommendations/>.

BACKGROUND: Section 4106(b) of the Affordable Care Act helps make prevention affordable and accessible for all Americans by requiring health plans to cover preventive care services and eliminate cost sharing.

ACTION: Payments for claims with services dates on or after April 1, 2018, identified in the attached **“Preventive Care Services Guidelines”** provides guidance that providers may reference for billing purposes. Refer to the Agency for Healthcare Research and Quality (AHRQ) “The Guide to Clinical Preventive Services - 2014 Recommendation of the U.S. Preventive Services Task Force” for detailed recommendation clinical summaries which can be found at: <https://www.ahrq.gov/professionals/clinicians-providers/guidelines-recommendations/guide/index.html>.

Modifier 33 - Preventive Care Service

When the primary purpose of the service is the delivery of an evidence-based service in accordance with a US Preventive Services Task Force A and B rating in effect and other preventive care services identified in preventive care services mandates (legislative or regulatory), the service may be identified by modifier 33 to the procedure.

If you have any questions concerning this Newsletter, please contact Molina Medicaid Solutions (formerly Unisys) Provider Services at 1-800-776-6334 or <http://www.njmmis.com>.

RETAIN THIS NEWSLETTER FOR FUTURE REFERENCE



State of New Jersey
Division of Medical Assistance and Health Services

PREVENTIVE CARE SERVICES GUIDELINES

USPSTF A and B Recommendations	Service Description	CPT Code	ICD-10 Codes	Codes Criteria
Abdominal Aortic Aneurysm Screening Men	The USPSTF recommends one-time screening for abdominal aortic aneurysm by ultrasonography in men ages 65 to 75 years who have ever smoked.	76700 76705 76706 76770 76775 G0389	Z87.891 F17.200 F17.201 F17.210 F17.211 F17.213 F17.218 F17.219 F17.290 F17.291 Z13.6	Must have a listed Procedure code AND a listed ICD-10 Code
Alcohol Misuse: Screening & Counseling	The USPSTF recommends that clinicians screen adults age 18 years or older for alcohol misuse and provide persons engaged in risky or hazardous drinking with brief behavioral counseling interventions to reduce alcohol misuse.	Screening G0442 99408 99409 Counseling G0443	Any ICD-10 codes	Preventive benefit with ANY ICD-10 Code
Aspirin Preventive Medication: Adults Aged 50 to 59 years with a ≥10% 10-year cardiovascular risk	The USPSTF recommends initiating low-dose aspirin use for the primary prevention of cardiovascular disease and colorectal cancer in adults aged 50 to 59 years who have a 10% or greater 10-year cardiovascular risk, are not at increased risk for bleeding, have a life expectancy of at least 10 years, and are willing to take low-dose aspirin daily for at least 10 years.	G0446 Aspirin as prescribed by primary care clinicians	Any ICD-10 codes	Preventive benefit with ANY ICD-10 Code
Bacteriuria Screening - Pregnant Women	The USPSTF recommends screening for asymptomatic bacteriuria with urine culture in pregnant women at 12 to 16 weeks' gestation or at the first prenatal visit, if later.	81000 87077 87081 87086 87088	Pregnancy Codes	Must have a listed Procedure code AND a listed ICD-10 Code
Blood pressure screening: adults	The USPSTF recommends screening for high blood pressure in adults aged 18 years or older. The USPSTF recommends obtaining measurements outside of the clinical setting for diagnostic confirmation before starting treatment.	93784 93786 93788 93790	R03.0	Must have a listed Procedure code AND/OR a listed ICD-10 Code
BRCA Risk Assessment and Genetic Counseling/Testing	The USPSTF recommends that primary care providers screen women who have family members with breast, ovarian, tubal, or peritoneal cancer with one of several screening tools designed to identify a family history that may be associated with an increased risk for potentially harmful mutations in breast cancer susceptibility genes (BRCA1 or BRCA2). Women with positive screening results should receive genetic counseling and, if indicated after counseling, BRCA testing.	Genetic Counseling 96040 99201 99202 99203 99204 99205 99211 99212 99213 99214 99215 99385 99386 99387 99395 99396 99397 Testing 81211-81217 81162	Z80.3 Z80.41 Z31.5 Z15.01 Z15.02	Must have a listed Procedure code AND a listed ICD-10 Code

USPSTF A and B Recommendations	Service Description	CPT Code	ICD-10 Codes	Codes Criteria
Breast Cancer Preventive Medications	The USPSTF recommends that clinicians engage in shared, informed decisionmaking with women who are at increased risk for breast cancer about medications to reduce their risk. For women who are at increased risk for breast cancer and at low risk for adverse medication effects, clinicians should offer to prescribe risk-reducing medications, such as tamoxifen or raloxifene.	99401 99402 99403 99404	Z80.3 Z80.41 Z15.01 Z15.02	Must have a listed Procedure Code AND/OR a listed ICD-10 Code
Breast Cancer Screening (Mammography)	The USPSTF recommends screening mammography for women, with or without clinical breast examination, every 1 to 2 years for women age 40 years and older.	G0202 77052 77055 77056 77057 77065 77066 77067	Preventive benefit with any diagnosis code	ANY listed CPT/HCPC code listed
Breastfeeding interventions	The USPSTF recommends providing interventions during pregnancy and after birth to support breastfeeding.	Support & Counseling 99241-99245 99341-99345 99347-99350 Equipment & Supplies E0602 E0603 E0604 A4281 A4282 A4283 A4284 A4285 A4286	Z39.1 P92.1 P92.2 P92.3 P92.4 P92.5 P92.8 P92.9 R63.3	Must have a listed Procedure Code AND a listed ICD-10 Code Preventive benefit with ANY diagnosis code
Cervical Cancer Screening (Pap Smear)	The USPSTF recommends screening for cervical cancer in women ages 21 to 65 years with cytology (Pap smear) every 3 years or, for women ages 30 to 65 years who want to lengthen the screening interval, screening with a combination of cytology and human papillomavirus (HPV) testing every 5 years.	G0101 G0123 G0124 G0141 Q0091 P3000 P3001 88141-88143 88147 88148 88150 88152-88155 88164-88167 88174 88175	Z00.00 Z00.01 Z01.411 Z01.419 Z01.42 Z12.4	Preventive benefit with ANY diagnosis code
HPV SCREENING		87623 87624 87625	Z11.51	Preventive benefit with ANY diagnosis code
Chlamydia Screening: Women	The USPSTF recommends screening for chlamydia in sexually active women age 24 years or younger and in older women who are at increased risk for infection.(including pregnant women).	87110 87270 87320 87490 87491 87492 87810 86631 86632	Z11.3 Z11.8 Z11.9 Z72.89 Pregnancy codes	Preventive benefit with ANY diagnosis code Must have a listed Procedure code AND an ICD 10 pregnancy code.

USPSTF A and B Recommendations	Service Description	CPT Code	ICD-10 Codes	Codes Criteria
Colorectal Cancer Screening	The USPSTF recommends screening for colorectal cancer starting at age 50 years and continuing until age 75 years.	Fecal Occult Blood Testing 81528 G0464 82270 82274 G0328 Sigmoidoscopy G0104 G0106 45330 45331 45333 45338 45346 Barium Enema G0122	Z80.0 Z83.71 Z12.12 Z12.11 Z83.79 Z00.00 Z00.01 Z12.10	Must have a listed Procedure Code AND a listed ICD-10 Code
Colorectal Cancer Screening	The USPSTF recommends screening for colorectal cancer starting at age 50 years and continuing until age 75 years.	Colonoscopy G0105 G0120 G0121 45378 45380 45384 45385 45388 44388 44389 44392 44401 44394 CT Scan 74261 74262 74263		Must have a listed Procedure Code AND a listed ICD-10 Code
Dental caries prevention: infants and children up to age 5 years	The USPSTF recommends the application of fluoride varnish to the primary teeth of all infants and children starting at the age of primary tooth eruption in primary care practices. The USPSTF recommends primary care clinicians prescribe oral fluoride supplementation starting at age 6 months for children whose water supply is fluoride deficient.	99188	Z29.3	Must have a listed Procedure Code AND/OR a listed ICD-10 Code
Depression Screening: Adolescent	The USPSTF recommends screening for major depressive disorder (MDD) in adolescents aged 12 to 18 years. Screening should be implemented with adequate systems in place to ensure accurate diagnosis, effective treatment, and appropriate follow-up.	G0444 96127	Z13.89	Must have a listed Procedure Code AND/OR a listed ICD-10 Code
Depression Screening: Adult	The USPSTF recommends screening for depression in the general adult population, including pregnant and postpartum women. Screening should be implemented with adequate systems in place to ensure accurate diagnosis, effective treatment, and appropriate follow-up.	G0444 96127	Z13.89	Must have a listed Procedure Code AND/OR a listed ICD-10 Code
Diabetes Screening	The USPSTF recommends screening for abnormal blood glucose as part of cardiovascular risk assessment in adults aged 40 to 70 years who are overweight or obese. Clinicians should offer or refer patients with abnormal blood glucose to intensive behavioral counseling interventions to promote a healthful diet and physical activity.	82947 82948 82950 82951 83036	Z00.00 Z00.01 Z13.1 Z13.6 Z13.29 Obesity Codes HTN Codes	Must have a listed Procedure Code AND a listed ICD-10 Code
Falls Prevention in Older Adults: Exercise or Physical Therapy	The USPSTF recommends exercise or physical therapy to prevent falls in community-dwelling adults age 65 years and older who are at increased risk for falls.	97110 97112 97116 97113 97150 97530	Z91.81 Z00.00 Z00.01 M62.81 W19.XXXS	Must have a listed Procedure Code AND a listed ICD-10 Code

USPSTF A and B Recommendations	Service Description	CPT Code	ICD-10 Codes	Codes Criteria
Falls Prevention in older Adults: Vitamin D	The USPSTF recommends vitamin D supplementation to prevent falls in community-dwelling adults age 65 years and older who are at increased risk for falls.	Vitamin D supplementation as prescribed by primary care clinicians	N/A	
Folic Acid Supplementation	The USPSTF recommends that all women who are planning or capable of pregnancy take a daily supplement containing 0.4 to 0.8 mg (400 to 800 µg) of folic acid.	Folic Acid Supplementation as prescribed by primary care clinicians	N/A	
Gestational Diabetes Mellitus Screening	The USPSTF recommends screening for gestational diabetes mellitus in asymptomatic pregnant women after 24 weeks of gestation.	82947 82950 82951 82952 83036	Pregnancy codes	Must have a listed Procedure Code AND a listed ICD-10 Code
Gonorrhea Prophylactic Medication: Newborns (Ocular prophylaxis for gonococcal neonatorum preventive medication)	The USPSTF recommends prophylactic ocular topical medication for all newborns for the prevention of gonococcal ophthalmia neonatorum.	99460 99461 99463 Newborn DRG 580-640 99391	Prophylactic Ocular Topical Medication administered at time of delivery Z00.110 Z29.9	Must have a listed Procedure Code AND a listed ICD-10 Code OR DRG
Gonorrhea Screening: Women	The USPSTF recommends screening for gonorrhea in sexually active women age 24 years or younger and in older women who are at increased risk for infection.	87081 87205 87590 87591 87592 87801 87850	Z72.51 Z72.52 Z72.53 Z72.89 Z11.3 Pregnancy Codes	Must have a listed Procedure Code AND a listed ICD-10 Code Must have pregnancy code AND procedure code
Healthy Diet and physical activity counseling to prevent cardiovascular disease: Adults with cardiovascular risk factors.	The USPSTF recommends offering or referring adults who are overweight or obese and have additional cardiovascular disease (CVD) risk factors to intensive behavioral counseling interventions to promote a healthful diet and physical activity for CVD prevention.	Behavior Counseling or Therapy G0446 G0447 G0473 Medical Nutrition Therapy or Counseling 97802 97803 Preventive Medicine Individual Counseling 99401 99402 99403 99404	E66.09 E66.1 E66.8 E66.9 E66.01 E66.3 E78.4 E78.5 Z13.6 Z71.3 Obesity Codes Tobacco Use Codes Diabetes Codes	Must have a listed Procedure Code AND a listed ICD-10 Code
Hemoglobinopathies Screening: Newborns	The USPSTF recommends screening for sickle cell disease in newborns.	S3620 83020 83021 85660	Z00.121 Z00.129 Z13.0	Must have a listed Procedure Code AND a listed ICD-10 Code or DRG
Hepatitis B Screening: Nonpregnant Adolescents and Adults	The USPSTF recommends screening for hepatitis B virus infection in persons at high risk for infection.	86704 - 86707 87340- 87341	Z20.5 Z20.828 Z11.59	Must have a listed Procedure Code AND a listed ICD-10 Code
Hepatitis B Screening: Pregnant Women	The USPSTF strongly recommends screening for hepatitis B virus infection in pregnant women at their first prenatal visit.	80055 86704 -86707 87340-87341	Pregnancy codes	Must have a listed Procedure Code AND a listed ICD-10 Code

USPSTF A and B Recommendations	Service Description	CPT Code	ICD-10 Codes	Codes Criteria
Screening for Hepatitis C Virus Infection: Adults	The USPSTF recommends screening for hepatitis C virus (HCV) infection in persons at high risk for infection. The USPSTF also recommends offering one-time screening for HCV infection to adults born between 1945 and 1965.	86803 86804 87522	F19.20 F19.21 Z99.2 Z51.89 Z72.51 Z72.52 Z72.53 Z72.89 B17.10 B18.2 B19.20 W46.1XXA W46.1XXD Z00.00 Z00.01 Z00.8 Z21 Z05.1 Hepatitis Codes	Must have a listed Procedure Code AND a listed ICD-10
HIV Screening: nonpregnant adolescents and adults	The USPSTF recommends that clinicians screen for HIV infection in adolescents and adults ages 15 to 65 years. Younger adolescents and older adults who are at increased risk should also be screened.	G0432 G0433 G0435 86689 86701 86702 86703 87534 87535 87536 87389 87390 87391	Z72.89 Z00.00 Z00.01 Z11.4 Z11.59	Must have a listed Procedure Code AND a listed ICD-10 Code
HIV Screening: pregnant women	The USPSTF recommends that clinicians screen all pregnant women for HIV, including those who present in labor who are untested and whose HIV status is unknown.	G0432 G0433 G0435 86689 86701 86702 86703 87534 87535 87536 87389 87390 87391	Pregnancy codes	Must have a listed Procedure Code AND a listed ICD-10 Code.
Hypothyroidism Screening: Newborns	The USPSTF recommends screening for congenital hypothyroidism in newborns.	99460 99461 99463 S3620	P09 Z00.110 Z13.29	Must have a listed DRG.Code AND a listed ICD-10 Code
Intimate Partner Violence Screening: Women of Childbearing Age	The USPSTF recommends that clinicians screen women of childbearing age for intimate partner violence, such as domestic violence, and provide or refer women who screen positive to intervention services. This recommendation applies to women who do not have signs or symptoms of abuse.	99384 99385 99401 99402 99403 99404	T74.91XA T76.91XA T74.11XA T76.11XA T74.31XA T76.31XA T74.21XA T76.21XA T74.01XA T76.01XA Y07.12 Z71.89 Z65.8	Must have a listed Procedure Code AND a listed ICD-10 Code.
Lung Cancer Screening	The USPSTF recommends annual screening for lung cancer with low-dose computed tomography in adults ages 55 to 80 years who have a 30 pack-year smoking history and currently smoke or have quit within the past 15 years. Screening should be discontinued once a person has not smoked for 15 years or develops a health problem that substantially limits life expectancy or the ability or willingness to have curative lung surgery.	G0296 G0297 71250 76497	Z87.891 Z12.2 Tobacco Use Codes	Must have a listed Procedure code AND a listed ICD-10 code

USPSTF A and B Recommendations	Service Description	CPT Code	ICD-10 Codes	Codes Criteria
Obesity screening and counseling: Adults	The USPSTF recommends screening all adults for obesity. Clinicians should offer or refer patients with a body mass index of 30 kg/m ² or higher to intensive, multicomponent behavioral interventions.	Preventive Medicine Individual Counseling 99401 99402 99403 99404 Medical Nutrition Therapy 97802 97803 Behavioral Counseling or Therapy G0446 G0447 G0473	Z68.30 Z68.31 Z68.32 Z68.33 Z68.34 Z68.35 Z68.36 Z68.37 Z68.38 Z68.39 Z68.41 Z68.42 Z68.43 Z68.44 Z68.45 E66.09 E66.1 E66.8 E66.9 E66.01 Obesity Codes	Must have a listed Procedure Code AND a listed ICD-10 Code
Obesity screening : Children and Adolescents	The USPSTF recommends that clinicians screen for obesity in children and adolescents 6 years and older and offer or refer them to comprehensive, intensive behavioral interventions to promote improvement in weight status.	Preventive Medicine Individual Counseling 99401 99402 99403 99404 Medical Nutrition Therapy 97802 97803 Behavioral Counseling or Therapy G0446 G0447 G0473	Z68.53 Z68.54 E66.09 E66.1 E66.8 E66.9 E66.01 Obesity Codes	Must have a listed Procedure Code AND a listed ICD-10 Code
Osteoporosis Screening: Women	The USPSTF recommends screening for osteoporosis in women age 65 years and older and in younger women whose fracture risk is equal to or greater than that of a 65-year-old white woman who has no additional risk factors.	77080 77081 G0130	Z13.820 Z78.0 M81.0 M81.6 M81.8 M85.9 M89.9	Must have a listed Procedure Code AND a listed ICD-10 Code
Phenylketonuria Screening: Newborns	The USPSTF recommends screening for phenylketonuria in newborns.	S3620 84030 99460 99461 99463	Z76.1 Z00.129 Z00.110 Z13.228	Must have listed Procedure Code AND a listed ICD-10 Code or DRG
Preeclampsia Prevention: Aspirin	The USPSTF recommends the use of low-dose aspirin (81 mg/d) as preventive medication after 12 weeks of gestation in women who are at high risk for preeclampsia.	Low-dose aspirin as prescribed by primary care clinicians	N/A	N/A
Preeclampsia Screening	The USPSTF recommends screening for preeclampsia in pregnant women with blood pressure measurements throughout pregnancy.	59400 59510 59610 59618	Pregnancy Codes	Must have a listed Procedure Code AND/OR a listed ICD-10 Code
RH Incompatibility Screening: first pregnancy visit	The USPSTF strongly recommends Rh (D) blood typing and antibody testing for all pregnant women during their first visit for pregnancy-related care.	80055 86900 86901	Pregnancy Codes O36.0110 O36.0120 O36.0130 O36.0910 O36.0920 O36.0930 Z01.83 Z31.7 Z31.89	Must have a listed Procedure Code AND a listed ICD-10 Code

USPSTF A and B Recommendations	Service Description	CPT Code	ICD-10 Codes	Codes Criteria
RH Incompatibility Screening: 24-28 weeks' gestation	The USPSTF recommends repeated Rh (D) antibody testing for all unsensitized Rh (D)-negative women at 24 to 28 weeks' gestation, unless the biological father is known to be Rh (D)-negative.	80055 86900 86901	Pregnancy Codes P07.23 P07.24 P07.25 P07.26 P07.31 P55.0 O36.0110 O36.0120 O36.0130 O36.0910 O36.0920 O36.0930 Z01.83 Z31.7 Z31.89	Must have a listed Procedure Code AND a listed ICD-10 Code
Sexually Transmitted Infections Counseling	The USPSTF recommends intensive behavioral counseling for all sexually active adolescents and for adults who are at increased risk for sexually transmitted infections.	G0445 99401 99402 99403 99404	Z70.1 Z72.5 Z72.51 Z72.52 Z72.53 Z72.89 Z11.3	Must have a listed Procedure Code AND a listed ICD-10 Code
Skin Cancer Behavioral Counseling	The USPSTF recommends counseling young adults, adolescents, children and parents of young children about minimizing exposure to ultraviolet (UV) radiation for persons aged 6 months to 24 years with fair skin types to reduce their risk of skin cancer.	99401 99402 99403 99404	Z12.83	Must have a listed Procedure Code AND a listed ICD-10 Code
Statin Preventive Medication: Adults age 40-75 years with no history of CVD, 1 or more CVD risk factors, and a calculated 10 year CVD event risk of 10% or greater.	The USPSTF recommends that adults without a history of cardiovascular disease (CVD) (i.e., symptomatic coronary artery disease or ischemic stroke) use a low- to moderate-dose statin for the prevention of CVD events and mortality when all of the following criteria are met: 1) they are ages 40 to 75 years; 2) they have 1 or more CVD risk factors (i.e., dyslipidemia, diabetes, hypertension, or smoking); and 3) they have a calculated 10-year risk of a cardiovascular event of 10% or greater. Identification of dyslipidemia and calculation of 10-year CVD event risk requires universal lipids screening in adults ages 40 to 75 years.	G9664	Hypertension Codes Tobacco Use Codes Diabetes Codes E78.0 E78.1 E78.2 E78.4 E78.5	Must have a listed Procedure Codes AND/OR a listed ICD-10 Code
Tobacco Use Counseling and Interventions: Non-Pregnant Adults	The USPSTF recommends that clinicians ask all adults about tobacco use, advise them to stop using tobacco, and provide behavioral interventions and U.S. Food and Drug Administration (FDA)-approved pharmacotherapy for cessation to adults who use tobacco.	99406 99407	Z72.0 Z87.891 F17.200 F17.201 F17.210 F17.211 F17.220 F17.221 F17.290 F17.291 T65.2-T65.294	Must have a listed Procedure Code AND a listed ICD-10 Code
Tobacco Use Counseling: Pregnant Women	The USPSTF recommends that clinicians ask all pregnant women about tobacco use, advise them to stop using tobacco, and provide behavioral interventions for cessation to pregnant women who use tobacco.	99406 99407	Pregnancy Codes Z72.0 Z87.891 O99.330 O99.331 O99.332 O99.333 O99.334 O99.335 T65.2 - T65.294	Must have a listed Procedure Code AND a listed ICD-10 Code

USPSTF A and B Recommendations	Service Description	CPT Code	ICD-10 Codes	Codes Criteria
Tobacco Use Interventions: Children and Adolescents	The USPSTF recommends that clinicians provide interventions, including education or brief counseling, to prevent initiation of tobacco use in school-aged children and adolescents.	99406 99407	Z72.0 Z87.891 F17.200 F17.201 F17.210 F17.211 F17.220 F17.221 F17.290 F17.291 Z87.891 Z77.22	Must have a listed Procedure Code AND a listed ICD-10 Code
Tuberculosis screening: adults	The USPSTF recommends screening for latent tuberculosis infection in populations at increased risk.	86480 86580	R76.11 R76.12 Z11.1	Must have a listed Procedure Code AND/OR a listed ICD-10 Code
Syphilis Screening: Non-pregnant Persons	The USPSTF recommends screening for syphilis infection in persons who are at increased risk for infection.	86592 86593 86780	Z00.121 Z00.129 Z72.89 Z00.0 Z00.01 Z11.2 Z11.9 Z20.2	Must have a listed Procedure Code AND a listed ICD-10 Code
Syphilis Screening: Pregnant Women	The USPSTF recommends that clinicians screen all pregnant women for syphilis infection.	80055 86592 86593 86780	Pregnancy Codes	Must have a listed Procedure Code AND a listed ICD-10 Code
Visual Screening: Children	The USPSTF recommends vision screening for all children at least once between the ages of 3 and 5 years, to detect the presence of amblyopia or its risk factors.	99173 99174 99177	Z76.2 Z00.121 Z00.129	Must have a listed Procedure Code AND a listed ICD-10 Code

ADULT VACCINES	
	CPT Code
INFLUENZA	
Fluzone Intradermal quadrivalent	90630
Fluad	90653
Fluzone Intradermal trivalent	90654
Afluria Trivalent	<i>Preservative Free</i>
Fluzone Trivalent	90656
FluLaval Trivalent	
Fluvirin Trivalent	
Fluarix Trivalent	
Afluria Trivalent	<i>Preservative</i>
FluLaval Trivalent	90658
Fluzone Trivalent	
Fluvirin Trivalent	
FluMist Trivalent (intranasal)	90660
Flucelvax Trivalent	90661
Fluzone High Dose	90662
FluMist Quadrivalent (intranasal)	90672
Flublok Trivalent	90673
Flucelvax Quadrivalent	90674
Afluria Quadrivalent	<i>Preservative Free</i>
Fluarix Quadrivalent	90686
Fluzone Quadrivalent	
Flulaval Quadrivalent	
Afluria Quadrivalent	<i>Preservative</i>
FluLaval Quadrivalent	90688
Fluzone Quadrivalent	
TETANUS, DIPHTHERIA, AND PERTUSSIS	
Adacel	90715
Boostrix	
TETANUS AND DIPHTHERIA	
Tenivac	90714
TD (generic)	
VARICELLA	
Varivax	90716
HUMAN PAPILLOMA VIRUS (HPV)	
Gardasil HPV 4	90649
Cervarix	90650
Gardasil 9	90651
ZOSTER/SHINGLES (HZV)	
Zostavax	90736
MEASLES, MUMPS, RUBELLA (MMR)	
M-M-R II	90707
PNEUMOCOCCAL 16-VALENT CONJUGATE (PCV13)	
Pevnar 13	90670
PNEUMOCOCCAL POLYSACCHARIDE (PPSV23)	
Pneumovax 23	90732
MENINGOCOCCAL	
Bexsero	90620
Trumenba	90621
Menomune	90733
Menactra	
Menveo	90734
HEPATITIS A	
Havrix	<i>Adult 2 Dose</i>
Vaqta	90632
HEPATITIS A & B	
TwinRix	90636
HEPATITIS B	
Recombivax HB	<i>Dialysis 3 Dose</i>
	90740
Recombivax HB	<i>Adult 3 Dose</i>
Energix B	90746
EnGerix-B	<i>Dialysis 4 Dose</i>
	90747
HEPATITIS B and HAEMOPHILUS INFLUENZA TYPE B	
Comvax	90748

ICD-10 ATHEROSCLEROSIS CODES

170.0	170.245	170.329	170.403	170.461	170.534	170.612	170.669	170.741
170.1	170.248	170.331	170.408	170.462	170.535	170.613	170.691	170.742
170.201	170.249	170.332	170.409	170.463	170.538	170.618	170.692	170.743
170.202	170.25	170.333	170.411	170.468	170.539	170.619	170.693	170.744
170.203	170.261	170.334	170.412	170.469	170.541	170.621	170.698	170.745
170.208	170.262	170.335	170.413	170.491	170.542	170.622	170.699	170.748
170.209	170.263	170.338	170.418	170.492	170.543	170.623	170.701	170.749
170.211	170.268	170.339	170.419	170.493	170.544	170.628	170.702	170.75
170.212	170.269	170.341	170.421	170.498	170.545	170.629	170.703	170.761
170.213	170.291	170.342	170.422	170.499	170.548	170.631	170.708	170.762
170.218	170.292	170.343	170.423	170.501	170.549	170.632	170.709	170.763
170.219	170.293	170.344	170.428	170.502	170.55	170.633	170.711	170.768
170.221	170.298	170.345	170.429	170.503	170.561	170.634	170.712	170.769
170.222	170.299	170.348	170.431	170.508	170.562	170.635	170.713	170.791
170.223	170.301	170.349	170.432	170.509	170.563	170.638	170.718	170.792
170.228	170.302	170.35	170.433	170.511	170.568	170.639	170.719	170.793
170.229	170.303	170.361	170.434	170.512	170.569	170.641	170.721	170.798
170.231	170.308	170.362	170.435	170.513	170.591	170.642	170.722	170.799
170.232	170.309	170.363	170.438	170.518	170.592	170.643	170.723	170.8
170.233	170.311	170.368	170.439	170.519	170.593	170.644	170.728	170.90
170.234	170.312	170.369	170.441	170.521	170.598	170.645	170.729	170.91
170.235	170.313	170.391	170.442	170.522	170.599	170.648	170.731	
170.238	170.318	170.392	170.443	170.523	170.601	170.649	170.732	
170.239	170.319	170.393	170.444	170.528	170.602	170.65	170.733	
170.241	170.321	170.398	170.445	170.529	170.603	170.661	170.734	
170.242	170.322	170.399	170.448	170.531	170.608	170.662	170.735	
170.243	170.323	170.401	170.449	170.532	170.609	170.663	170.738	
170.244	170.328	170.402	170.45	170.533	170.611	170.668	170.739	

ICD-10 DIABETES CODES

E08.3211	E08.3532	E09.3393	E09.3599	E10.3521	E11.3312	E11.3553	E13.3499
E08.3212	E08.3533	E09.3399	E09.37X1	E10.3522	E11.3313	E11.3559	E13.3511
E08.3213	E08.3539	E09.3411	E09.37X2	E10.3523	E11.3319	E11.3591	E13.3512
E08.3219	E08.3541	E09.3412	E09.37X3	E10.3529	E11.3391	E11.3592	E13.3513
E08.3291	E08.3542	E09.3413	E09.37X9	E10.3531	E11.3392	E11.3593	E13.3519
E08.3292	E08.3543	E09.3419	E10.3211	E10.3532	E11.3393	E11.3599	E13.3521
E08.3293	E08.3549	E09.3491	E10.3212	E10.3533	E11.3399	E11.37X1	E13.3522
E08.3299	E08.3551	E09.3492	E10.3213	E10.3539	E11.3411	E11.37X2	E13.3523
E08.3311	E08.3552	E09.3493	E10.3219	E10.3541	E11.3412	E11.37X3	E13.3529
E08.3312	E08.3553	E09.3499	E10.3291	E10.3542	E11.3413	E11.37X9	E13.3531
E08.3313	E08.3559	E09.3511	E10.3292	E10.3543	E11.3419	E13.3211	E13.3532
E08.3319	E08.3591	E09.3512	E10.3293	E10.3549	E11.3491	E13.3212	E13.3533
E08.3391	E08.3592	E09.3513	E10.3299	E10.3551	E11.3492	E13.3213	E13.3539
E08.3392	E08.3593	E09.3519	E10.3311	E10.3552	E11.3493	E13.3219	E13.3541
E08.3393	E08.3599	E09.3521	E10.3312	E10.3553	E11.3499	E13.3291	E13.3542
E08.3399	E08.37X1	E09.3522	E10.3313	E10.3559	E11.3511	E13.3292	E13.3543
E08.3411	E08.37X2	E09.3523	E10.3319	E10.3591	E11.3512	E13.3293	E13.3549
E08.3412	E08.37X3	E09.3529	E10.3391	E10.3592	E11.3513	E13.3299	E13.3551
E08.3413	E08.37X9	E09.3531	E10.3392	E10.3593	E11.3519	E13.3311	E13.3552
E08.3419	E09.3211	E09.3532	E10.3393	E10.3599	E11.3521	E13.3312	E13.3553
E08.3491	E09.3212	E09.3533	E10.3399	E10.37X1	E11.3522	E13.3313	E13.3559
E08.3492	E09.3213	E09.3539	E10.3411	E10.37X2	E11.3523	E13.3319	E13.3591
E08.3493	E09.3219	E09.3541	E10.3412	E10.37X3	E11.3529	E13.3391	E13.3592
E08.3499	E09.3291	E09.3542	E10.3413	E10.37X9	E11.3531	E13.3392	E13.3593
E08.3511	E09.3292	E09.3543	E10.3419	E11.3211	E11.3532	E13.3393	E13.3599
E08.3512	E09.3293	E09.3549	E10.3491	E11.3212	E11.3533	E13.3399	E13.37X1
E08.3513	E09.3299	E09.3551	E10.3492	E11.3213	E11.3539	E13.3411	E13.37X2
E08.3519	E09.3311	E09.3552	E10.3493	E11.3219	E11.3541	E13.3412	E13.37X3
E08.3521	E09.3312	E09.3553	E10.3499	E11.3291	E11.3542	E13.3413	E13.37X9
E08.3522	E09.3313	E09.3559	E10.3511	E11.3292	E11.3543	E13.3419	
E08.3523	E09.3319	E09.3591	E10.3512	E11.3293	E11.3549	E13.3491	
E08.3529	E09.3391	E09.3592	E10.3513	E11.3299	E11.3551	E13.3492	
E08.3531	E09.3392	E09.3593	E10.3519	E11.3311	E11.3552	E13.3493	

ICD-10 HEPATITIS C CODES

A50.01	A52.71	A60.01	F13.232	F18.251	K51.313	N49.9	Z20.2
A50.02	A52.72	A60.02	F13.239	F18.259	K51.314	N73.5	Z20.5
A50.03	A52.73	A60.03	F13.24	F18.27	K51.318	N73.9	Z20.6
A50.04	A52.74	A60.04	F13.250	F18.280	K51.319	N76.0	Z20.828
A50.05	A52.75	A60.09	F13.251	F18.288	K51.40	N76.1	Z21
A50.06	A52.76	A60.1	F13.259	F18.29	K51.411	N76.2	Z22.4
A50.07	A52.77	A60.9	F13.26	F19.20	K51.412	N76.3	Z41.8
A50.08	A52.78	A63.0	F13.27	F19.21	K51.413	N77.1	Z48.21
A50.09	A52.79	A63.8	F13.280	F19.220	K51.414	O35.3XX0	Z48.22
A50.1	A52.8	A64	F13.281	F19.221	K51.418	O35.3XX1	Z48.24
A50.2	A52.9	A74.81	F13.282	F19.222	K51.419	O35.3XX2	Z48.280
A50.30	A53.0	A74.89	F13.288	F19.229	K51.50	O35.3XX3	Z48.288
A50.31	A53.9	A74.9	F13.29	F19.230	K51.511	O35.3XX4	Z48.290
A50.32	A54.00	B07.8	F14.20	F19.231	K51.512	O35.3XX5	Z48.298
A50.39	A54.01	B07.9	F14.21	F19.232	K51.513	O35.3XX9	Z49.31
A50.40	A54.02	B20.	F14.220	F19.239	K51.514	O35.5XX0	Z49.32
A50.41	A54.03	B97.35	F14.221	F19.24	K51.518	O35.5XX1	Z51.89
A50.42	A54.09	B97.7	F14.222	F19.250	K51.519	O35.5XX2	Z52.000
A50.43	A54.1	D65	F14.229	F19.251	K51.80	O35.5XX3	Z52.001
A50.44	A54.21	D66	F14.23	F19.259	K51.811	O35.5XX4	Z52.008
A50.45	A54.22	D67.	F14.24	F19.26	K51.812	O35.5XX5	Z52.010
A50.49	A54.23	D68.0	F14.250	F19.27	K51.813	O35.5XX9	Z52.011
A50.51	A54.24	D68.1	F14.251	F19.280	K51.814	O90.4	Z52.018
A50.52	A54.29	D68.2	F14.259	F19.281	K51.818	O98.011	Z52.090
A50.53	A54.30	D68.311	F14.280	F19.282	K51.819	O98.012	Z52.091
A50.54	A54.31	D68.312	F14.281	F19.288	K51.90	O98.013	Z52.098
A50.55	A54.32	D68.318	F14.282	F19.29	K51.911	O98.019	Z52.10
A50.56	A54.33	D68.32	F14.288	K50.00	K51.912	O98.02	Z52.11
A50.57	A54.39	D68.4	F14.29	K50.011	K51.913	O98.03	Z52.19
A50.59	A54.40	D68.8	F15.20	K50.012	K51.914	O98.111	Z52.20
A50.6	A54.41	D68.9	F15.21	K50.013	K51.918	O98.112	Z52.21
A50.7	A54.42	F11.20	F15.220	K50.014	K51.919	O98.113	Z52.29
A50.9	A54.43	F11.21	F15.221	K50.018	M02.30	O98.119	Z52.3
A51.0	A54.49	F11.220	F15.222	K50.019	M02.311	O98.12	Z52.4
A51.1	A54.5	F11.221	F15.229	K50.10	M02.312	O98.13	Z52.5

ICD-10 HEPATITIS C CODES

A51.2	A54.6	F11.222	F15.23	K50.111	M02.319	O98.211	Z52.6
A51.31	A54.81	F11.229	F15.24	K50.112	M02.321	O98.212	Z52.89
A51.32	A54.82	F11.23	F15.250	K50.113	M02.322	O98.213	Z52.9
A51.39	A54.83	F11.24	F15.251	K50.114	M02.329	O98.219	Z57.8
A51.41	A54.84	F11.250	F15.259	K50.118	M02.331	O98.22	Z71.7
A51.42	A54.85	F11.251	F15.280	K50.119	M02.332	O98.23	Z72.51
A51.43	A54.86	F11.259	F15.281	K50.80	M02.339	O98.311	Z72.52
A51.44	A54.89	F11.281	F15.282	K50.811	M02.341	O98.312	Z72.53
A51.46	A54.9	F11.282	F15.288	K50.812	M02.342	O98.313	Z72.9
A51.49	A55	F11.288	F15.29	K50.813	M02.349	O98.319	Z86.2
A51.5	A56.00	F11.29	F16.20	K50.814	M02.351	O98.32	Z92.25
A51.9	A56.01	F12.20	F16.21	K50.818	M02.352	O98.33	Z94.0
A52.00	A56.02	F12.21	F16.220	K50.819	M02.359	O99.320	Z94.1
A52.01	A56.09	F12.220	F16.221	K50.90	M02.361	O99.321	Z94.2
A52.02	A56.11	F12.221	F16.229	K50.911	M02.362	O99.322	Z94.3
A52.03	A56.19	F12.222	F16.24	K50.912	M02.369	O99.323	Z94.5
A52.04	A56.2	F12.229	F16.250	K50.913	M02.371	O99.324	Z94.6
A52.05	A56.3	F12.250	F16.251	K50.914	M02.372	O99.325	Z94.7
A52.06	A56.4	F12.251	F16.259	K50.918	M02.379	P00.2	Z94.81
A52.09	A56.8	F12.259	F16.280	K50.919	M02.38	Z00.00	Z94.82
A52.10	A57	F12.280	F16.283	K51.20	M02.39	Z00.01	Z94.83
A52.11	A58	F12.288	F16.288	K51.211	N18.3	Z04.41	Z94.84
A52.12	A59.00	F12.29	F16.29	K51.212	N18.4	Z04.42	Z94.89
A52.13	A59.01	F13.20	F18.20	K51.213	N18.5	Z05.1	Z94.9
A52.14	A59.02	F13.21	F18.21	K51.214	N18.6	Z11.3	Z95.3
A52.15	A59.03	F13.220	F18.220	K51.218	N34.1	Z11.4	Z95.4
A52.16	A59.09	F13.221	F18.221	K51.219	N49.1	Z11.59	Z99.2
A52.17	A59.8	F13.229	F18.229	K51.30	N49.2	Z11.9	
A52.19	A59.9	F13.230	F18.24	K51.311	N49.3	Z14.01	
A52.2	A60.00	F13.231	F18.250	K51.312	N49.8	Z14.02	
A52.3							

ICD-10 HYPERTENSION CODES

ABNORMAL BLOOD PRESSURE READING WITHOUT DIAGNOSIS OF HYPERTENSION

R03.0

HYPERTENSION HEART DISEASE

I11.0

I11.9

HYPERTENSION CHRONIC KIDNEY DISEASE

I12.0

I12.9

HYPERTENSIVE HEART AND CHRONIC KIDNEY DISEASE

I13.0

I13.10

I13.11

I13.2

ESSENTIAL HYPERTENSION

I10

SECONDARY HYPERTENSION

I15.0

I15.1

I15.2

I15.8

I15.9

N26.2

HYPERTENSION COMPLICATING PREGNANCY CHILDBIRTH AND THE PUERPERIUM

O10.011 O10.113 O10.22 O10.411 O10.913 O11.5 O16.1

O10.012 O10.119 O10.23 O10.412 O10.919 O11.9 O16.2

O10.013 O10.12 O10.311 O10.413 O10.92 O13.1 O16.3

O10.019 O10.13 O10.312 O10.419 O10.93 O13.2 O16.4

O10.02 O10.211 O10.313 O10.42 O11.1 O13.3 O16.5

O10.03 O10.212 O10.319 O10.43 O11.2 O13.4 O16.9

O10.111 O10.213 O10.32 O10.911 O11.3 O13.5

O10.112 O10.219 O10.33 O10.912 O11.4 O13.9

URGENT/EMERGENCY/CRISIS HYPERTENSION

I16.0

I16.1

I16.9

ICD-10 OBESITY CODES

OVERWEIGHT

E66.3	Z68.27
Z68.25	Z68.28
Z68.26	Z68.29

BODY MASS INDEX 30.0 – 39.9

Z68.30	Z68.35
Z68.31	Z68.36
Z68.32	Z68.37
Z68.33	Z68.38
Z68.34	Z68.39

BODY MASS INDEX 40.0 AND OVER

Z68.41	Z68.44
Z68.42	Z68.45
Z68.43	

OBESITY:

E66.01	E66.8
E66.09	E66.9
E66.1	

ICD-10 PREGNANCY CODES

Z34.00	009.213	009.813	O30.009	O30.129	O30.819	Z3A.25
Z34.01	009.219	009.819	O30.011	O30.191	O30.821	Z3A.26
Z34.02	009.291	009.821	O30.012	O30.192	O30.822	Z3A.27
Z34.03	009.292	009.822	O30.013	O30.193	O30.823	Z3A.28
Z34.80	009.293	009.823	O30.019	O30.199	O30.829	Z3A.29
Z34.81	009.299	009.829	O30.031	O30.201	O30.891	Z3A.30
Z34.82	009.30	O36.80X0	O30.032	O30.202	O30.892	Z3A.31
Z34.83	009.31	O36.80X1	O30.033	O30.203	O30.893	Z3A.32
Z34.90	009.32	O36.80X2	O30.039	O30.209	O30.899	Z3A.33
Z34.91	009.33	O36.80X3	O30.041	O30.211	Z3A.00	Z3A.34
Z34.92	009.511	O36.80X4	O30.042	O30.212	Z3A.01	Z3A.35
Z34.93	009.512	O36.80X5	O30.043	O30.213	Z3A.08	Z3A.36
Z33.1	009.513	O36.80X9	O30.049	O30.219	Z3A.09	Z3A.37
009.00	009.519	009.70	O30.091	O30.221	Z3A.10	Z3A.38
009.01	009.521	009.71	O30.092	O30.222	Z3A.11	Z3A.39
009.02	009.522	009.72	O30.093	O30.223	Z3A.12	Z3A.40
009.03	009.523	009.73	O30.099	O30.229	Z3A.13	Z3A.41
009.10	009.529	009.891	O30.101	O30.291	Z3A.14	Z3A.42
009.11	009.611	009.892	O30.102	O30.292	Z3A.15	Z3A.49
009.12	009.612	009.893	O30.103	O30.293	Z3A.16	
009.13	009.613	009.899	O30.109	O30.299	Z3A.17	
009.291	009.619	009.90	O30.111	O30.801	Z3A.18	
009.40	009.621	009.91	O30.112	O30.802	Z3A.19	
009.41	009.622	009.92	O30.113	O30.803	Z3A.20	
009.42	009.623	009.93	O30.119	O30.809	Z3A.21	
009.43	009.629	O30.001	O30.121	O30.811	Z3A.22	
009.211	009.811	O30.002	O30.122	O30.812	Z3A.23	
009.212	009.812	O30.003	O30.123	O30.813	Z3A.24	

ICD-10 TOBACCO USE CODES

Z87.891	O9.330
Z72.00	O99.330
F17.200	O99.331
F17.201	O99.332
F17.210	O99.333
F17.211	O99.334
F17.220	O99.335
F17.221	
F17.290	
F17.291	
Z77.22	
T65.221A	